

Truncation exercises

Outline

SL_2

Groups of higher rank

A simple model for Eisenstein series

Truncation for convex polyhedra

Bounded polyhedra

Cones

A local version

Conclusion of the proof

Langlands' combinatorial lemma

The Maass-Selberg formula

Back to Arthur's truncation

This talk can be found at

<http://www.math.ubc.ca/~cass/arthur/talk.pdf>


To start

$$\Gamma = \mathrm{SL}_2(\mathbb{Z})$$

$\mathcal{H} =$ **upper half plane**

$P =$ **Borel subgroup of upper triangular matrices**

$N =$ **unipotent matrices in P**



Thus for $Y \geq 1$ the quotient $\Gamma \cap P \setminus \mathcal{H}_Y$ may be identified with a subset of $\Gamma \setminus \mathcal{H}$. We have the map

$$\Gamma \cap P \setminus \mathcal{H}_Y \longrightarrow (0, \infty): z = x + iy \longmapsto y .$$

If χ_Y is the characteristic function of the region $y > Y$ truncation at Y is the operator


$$\Lambda^Y F = F - \chi_Y F_0$$

where

$$F_0(y) = \int_0^1 F(x + iy) dx$$

is the constant term of F .

The quotient $\Gamma \backslash \mathcal{H}$ may be compactified by adding a cusp at infinity, and truncation chops away the constant term of a function in the neighbourhood of the cusp.



As a Γ -invariant function on \mathcal{H}

$$\Lambda^Y F(z) = F(z) - \sum_{\Gamma \cap P \setminus \Gamma} \chi(y) F_0(y(\gamma z)) ,$$

- *Under a mild growth condition on F its truncation $\Lambda^Y F$ is rapidly decreasing at ∞ .*

Truncation plays a role in the meromorphic continuation of Eisenstein series and in proving the Selberg trace formula.

- *If E_s is the Eisenstein series of Maass then $\Lambda^Y E_s$ is square-integrable.*

The Maass-Selberg formula specifies $\|\Lambda^Y E_s\|^2$.

These features occur in using the truncation operator for groups of higher rank as well.

The Maass-Selberg formula:

$$\begin{aligned}\langle \Lambda^Y E_s, E_{-t} \rangle &= \langle \Lambda^Y E_s, E_{-t} \rangle \\ &= \int_0^Y (\text{const. term of } E_s)(\text{const. term of } E_{1-t}) \frac{dy}{y^2} \\ &= \int_0^Y (y^s + c(s)y^{1-s})(y^{1-t} + c(1-t)y^t) \frac{dx dy}{y^2}\end{aligned}$$

Groups of higher rank

Suppose now that G be a split group over \mathbb{Q} , $\Gamma = G(\mathbb{Z})$, $X = G/K$. All Borel subgroups are Γ -conjugate.

Fix one, call it P_\emptyset . Let Σ be the corresponding set of roots, Δ the basic roots.

For any rational parabolic subgroup P let N_P be its unipotent radical, $M_P = P/N_P$, A_P the connected component of the centre of M_P .

Given the compact subgroup K of G , for any P there exists a unique copy of M_P in P stable under the Cartan involution determined by K .

The $P \subseteq P_\emptyset$ are parametrized by subsets $\Theta \subseteq \Delta$.

For $\Theta \subseteq \Xi$

$$A_\Xi \subseteq A_\Theta \subseteq M_\Theta \subseteq M_\Xi$$

Every rational parabolic subgroup is Γ -conjugate to exactly one of these.

For any rational parabolic subgroup P and F on $\Gamma \backslash X$ the constant term of F with respect to P is


$$F_P(x) = \int_{\Gamma \cap N_P \backslash N_P} f(nx) \, dn ,$$

a function on $N_P(\Gamma \cap P) \backslash X$.

Conversely, for F on $N_P(\Gamma \cap P) \backslash X$ define (formally) the Eisenstein series

$$(E_P^G F)(x) = \sum_{\Gamma \cap P \backslash \Gamma} F(\gamma x) .$$

In A_P lie two naturally defined cones, one obtuse and one acute. Let χ_P be the characteristic function of the obtuse one, τ_P that of the acute one.



There is a canonical projection

$$N_P(\Gamma \cap P) \backslash X = N_P(\Gamma \cap P) \backslash P/K \cap P \longrightarrow A_P .$$

Let χ_P, τ_P be also their lifts back to X , $\chi_{P,p}$ and $\tau_{P,p}$ their shifts by p in P .

Fix T in the positive Weyl chamber in A_\emptyset far away from the walls. Arthur's definition of truncation is this:

$$\Lambda_G^T F = \sum_P (-1)^{\dim A_P - \dim A_G} E_P^G(\chi_{P,T} \cdot F_P)$$

The sum evaluated at any given element is finite.

Again:

$$\Lambda_G^T F = \sum_P (-1)^{\dim A_P - \dim A_G} E_P^G(\chi_{P,T} \cdot F_P)$$

If you are familiar with the geometry of $\Gamma \backslash X$ you will likely find this definition puzzling, because there is no longer any obvious relationship between truncation and the geometry of a compactification of X .

For groups of rational rank greater than one, Arthur's truncation is not local on any Satake compactification.

Nonetheless, there is no doubt that Arthur's definition is the correct one.

Under a mild growth condition on F the truncation $\Lambda^T F$ is rapidly decreasing at infinity.

Truncation is a projection operator, too.

It does not affect functions whose constant term support lies inside a well defined compact subset of $\Gamma \backslash X$. In particular it does not affect cusp forms.

Truncation is defined on every M_P as well as G itself. There is an equivalent recursive definition of truncation that defines it for G in terms of truncation on the other M_P .

Theorem. *We have an orthogonal decomposition*

$$F = \sum_P E_P^G(\tau_{P,T} \cdot \Lambda_{M_P}^T F_P) .$$

This is proven by means of a purely geometric lemma about obtuse simplicial cones, originally due to Langlands.

The rest of this talk will try to explain why the definition of truncation is reasonable, and why this theorem holds. *Without, however, proving either of them!*


A simple model for Eisenstein series

When trying to understand Arthur's calculations, I find it helpful to see what's going on in a much simpler situation, one where combinatorial difficulties are isolated from analytical ones.

Suppose given a split algebraic torus A and a root system Σ associated to it.

Let G be the algebraic group generated by the torus and the Weyl group W of Σ . In some sense, this is a reductive group in which the unipotent groups are reduced to shadows.

The parabolic subgroups in this scheme are parametrized by the faces of Weyl chambers. Given a face F , the associated parabolic of rational points is the subgroup generated by the torus and the subgroup W_P of W whose elements fix the points on the face.



The group Γ is just W .


The space X may be identified with the quotient of A by its torsion subgroup, which I will identify with the vector space V in which the (co)roots live.

Automorphic functions are the characters of V that are W -invariant, and the analogue of an Eisenstein series is the finite sum

$$(E_P^G F)(v) = \sum_{W_P \backslash W} F(wv)$$


that maps a function in V^{W_P} to one in V^W .

The natural definition of truncation associated to a point T in V is multiplication of a function F by the characteristic function of the convex hull of T .




If T is non-singular, then the faces of its convex hull \mathfrak{C}_T are parametrized by the ‘parabolic subgroups’.

The orthogonal decomposition is that corresponding to the partition of V according to the *nearest face* of the convex set \mathcal{C}_T .



That this agrees with Arthur's definition is not obvious.



The agreement of the two definitions is actually a special case of a much more general result about convex polyhedra.


This model is pretty simple, but it is not irrelevant to the ‘adult’. Among other things, some of its combinatorial geometry leads to a direct calculation of the Plancherel measure on the continuous spectrum.

Truncation for convex polyhedra

Let C be a closed convex polyhedron, $C^\circ \neq \emptyset$.



- If F is a face of codimension one of C , let E_F^C be the open exterior half-plane bounded by F .




- If F is any other proper face

$$E_F^C = \bigcap_{F \subseteq F_* \subseteq C} E_{F_*}^C$$

- If $F = C$ let $E_F^C = V$.

If $C^\circ = \emptyset$, replace it by its product with the vector space perpendicular to it in these definitions.



I do not assume C bounded.

Let \mathcal{E}_F^C be the characteristic function of E_F^C .

The following theorem is due to Ishida for cones, Brion and Vergne for nondegenerate bounded polyhedra. The case of cones was rediscovered by Kottwitz, who put it in an appendix to a paper written with Goresky and MacPherson. The result for arbitrary convex polyhedra seems to be new, even in two dimensions.


Theorem C. *We have*

$$\sum_{F \preceq C} (-1)^{\text{codim} F} \mathcal{E}_F^C = \text{char}_C .$$


Minkowski or Weyl might have discovered it.

A few simple cases:

- **Coordinate octant or simplicial cone**



- **Simplex**



Another formulation:

For any P in V

$$\sum_{P \in E_F^C} (-1)^{\text{codim} F} = \begin{cases} 1 & P \in C \\ 0 & \text{otherwise.} \end{cases}$$

This suggests that cohomological Euler-Poincaré characteristics are going to play a role:


$$\sum_{F \preceq C} (-1)^{\text{codim} F} = \begin{cases} (-1)^{\dim C} & \text{if } C \text{ is closed and bounded} \\ 0 & \text{if } C \text{ is a closed cone} \\ 1 & \text{if } C \text{ is open} \end{cases}$$

The first two are equivalent, since a suitable slice through a cone is a bounded polyhedron.

Two faces are allowed to have the same affine support. In effect, the Theorem is about a cell decomposition of C compatible with its convex structure. If a number of faces F_* partition an open geometric face F° , their total contribution is just

$$\sum_{F_* \subseteq F^\circ} (-1)^{\text{codim} F_*} E_{F_*}^C = E_F^C$$

since all $E_{F_*}^C = E_F^C$ here and the Euler-Poincaré characteristic of the open face is 1.




What's going on?

Bounded polyhedra *following Brion & Vergne*

We want to prove that for any point P

$$\sum_{P \in E_F^C} (-1)^{\text{codim} F} = \begin{cases} 1 & P \in C \\ 0 & \text{otherwise} \end{cases}$$



If P is a point of C this is immediate. If P is not in C , let H be the convex hull of C and P .



Then H is the union of all segments $[P, Q]$ with Q in C , and H° is the union of all $[Q, P)$ with Q in C° .

Proposition. *If F is a face of C then $F \subset H^\circ$ if and only if $P \in E_F^C$.*

Corollary. *A face F of C is one of the cells in the boundary of H if and only if $P \notin E_F^C$.*



$$\sum_{F \preceq H} (-1)^{\text{codim} F} = EP(H) = (-1)^{\dim C}$$

$$\sum_{F \preceq H, P \in F} (-1)^{\text{codim} F} = EP(\mathbf{cone}) = 0$$


$$\sum_{F \preceq H, P \notin F} (-1)^{\text{codim} F} = \sum_{F \preceq C, P \notin E_F^C} (-1)^{\text{codim} F} = (-1)^{\dim C}$$

$$\sum_{F \preceq C} (-1)^{\text{codim} F} = EP(C) = (-1)^{\dim C}$$

$$\sum_{P \in E_F^C} (-1)^{\text{codim} F} = 0 \quad \text{Q.E.D.}$$

Cones


The formula for cones reduces to that for bounded convex sets by taking slices.



Above the slice, the two configurations are the same.

A local version


For each face F of C , let V_F^C be the set of points in V for which the point of C nearest to it lies in F° .
Well, not quite.



It possesses an obvious
 product structure $F^\circ \times T_F^C$.

For each couple of faces $F_* \preceq F$ let

$$E_{F,F_*}^C = E_F^C \cap V_{F_*}^F.$$




Thus a point v of E_F^C lies in E_{F,F_*}^C if and only if the point of F closest to it lies in F_* .

Theorem L. *For each face F_* of C*





$$\sum_{F|F_* \preceq F} (-1)^{\operatorname{codim} F} \mathcal{E}_{F,F_*}^C = \begin{cases} 0 & F_* \neq C \\ \chi_C & F_* = C \end{cases}$$

This is one variation of Langlands' combinatorial lemma.

L in two dimensions is covered by these images . . .



... whose secret is given away by these:



C for cones implies L.

Theorem L asserts that

$$\sum_{F|F_* \preceq F} (-1)^{\text{codim } F} \mathcal{E}_{F,F_*}^C = \begin{cases} 0 & F_* \neq C \\ \chi_C & F_* = C \end{cases}$$

In this, C may be replaced by its tangent cone at F_* . At any face but a vertex, the tangent cone at that face has a simple product structure, and induction proves the claim. The formula for the full cone can be rearranged to give it for the vertex.





Conclusion of the proof of C

C follows from L by introducing the partition

$$\mathcal{E}_F^C = \sum_{F_* \preceq F} \mathcal{E}_{F, F_*}^C$$

and then rearranging the sum:

$$\begin{aligned} \sum_F (-1)^{\text{codim } F} \mathcal{E}_F^C &= \sum_{F, F_* \mid F_* \preceq F} (-1)^{\text{codim } F} \mathcal{E}_{F, F_*}^C \\ &= \sum_{F_*} \sum_{F \mid F_* \preceq F} (-1)^{\text{codim } F} \mathcal{E}_{F, F_*}^C \\ &= \chi_C . \end{aligned}$$



Langlands' combinatorial lemma

If F is a face of C , let T_F^C be the translation of V_F^C by the support of F , and τ_F^C its characteristic function. Thus $V_F^C = T_F^C \times F^\circ$. When C is an obtuse simplicial cone the following result is essentially the same as the original combinatorial lemma of Langlands.

Theorem. *For any face F of C*

$$\sum_{F \preceq F_* \preceq C} (-1)^{\text{codim} F} \tau_{F_*}^C \mathcal{E}_F^{F_*} = \begin{cases} 1 & \text{if } F = C \\ 0 & \text{otherwise} \end{cases}$$

The case $F = C$ is trivial. The proof for other F uses the partition of V into the V_F^C , and goes by induction.

The original applied to simplicial cones and was announced by Langlands without proof in his 1965 Boulder talk on Eisenstein series, and a result equivalent to this one is contained in the appendices to a recent paper by Goresky et al.

The Maass-Selberg formula

The Maass-Selberg formula is an explicit formula for the inner product of two Eisenstein series. Its analogue here is a formula for the Fourier transform of the characteristic function of a bounded convex set with $C^\circ \neq \emptyset$, observed by Brion & Vergne:

$$\widehat{\chi}_C(s) = (-1)^{\dim C} \sum_P \widehat{\mathcal{E}}_P^C(s)$$

where the right hand sum is over the vertices of C , and the expression is taken to be the analytic continuation of the obvious integral.

Applying the fundamental theorem to the exterior cones, this can be rewritten

$$\hat{\chi}_C(s) = \sum_P \hat{\mathcal{I}}_P^C(s)$$

where \mathcal{I} is the characteristic function of the *interiors* (tangent cones) of the vertices.

These integrals are easy to compute when these cones are simplicial, but as far as I know there is no simple formula otherwise.

Back to Arthur's truncation

I believe that Arthur's truncation is a generalization of this theory to the cone over the Tits building of the rational group G (Arthur, Stuhler, Grayson, Saper, Leuzinger, Ji & MacPherson). And that much of reduction theory is captured by this approach. Not quite clear yet.

References

J. Arthur, A trace formula for reductive groups I. Terms associated to classes in $G(\mathbb{Q})$, *Duke Mathematics Journal* 45 (1978), 911–952.

—————, A trace formula for reductive groups II. Applications of a truncation operator, *Compositio Mathematica* 40 (1980), 87–121.

M. Brion and M. Vergne, Lattice points in simple polytopes, *Journal of the American Mathematical Society* 10 (1997), 371–392.

M. Goresky, R. Kottwitz, and R. MacPherson, Discrete series characters and the Lefschetz formula for Hecke operators, *Duke Mathematics Journal* 89

(1997), 477–554. *Appendix B is the first place I am aware of where Langlands' combinatorial lemma is formulated for general cones.*

M. N. Ishida, Polyhedral Laurent series and Brion's inequalities, *International Journal of Mathematics* 1 (1990), 251–265.

J-P. Labesse, La formules de trace d'Arthur-Selberg, *Séminaire Bourbaki*, 1984–85, exposé 636.

R. P. Langlands, Some lemmas to be applied to the Eisenstein series, personal notes from around 1965 available at

<http://sunsite.ubc.ca/scans/lemma/cl.html>

—————, **Eisenstein series**, *Proceedings of Symposia in Pure Mathematics IX, 1965*. This was the Boulder conference. The relevant section is picturesquely called ' L^2 as the bed of Procrustes'.